

12-30-2019

Description of 2-local two sided multiplication on an algebra of matrices

Farhodjon N. Arzikulov

Andijan State University, Andijan, 170100, str. University, 129 (Uzbekistan). E-mail: agsu_info@edu.uz

Kamola A. Solijanov

Andijan State University, Andijan, 170100, str. University, 129 (Uzbekistan). E-mail: agsu_info@edu.uz

Follow this and additional works at: <https://uzjournals.edu.uz/adu>



Part of the [Mathematics Commons](#)

Recommended Citation

Arzikulov, Farhodjon N. and Solijanov, Kamola A. (2019) "Description of 2-local two sided multiplication on an algebra of matrices," *Scientific Bulletin. Physical and Mathematical Research*: Vol. 1 : No. 2 , Article 9.

Available at: <https://uzjournals.edu.uz/adu/vol1/iss2/9>

This Article is brought to you for free and open access by 2030 Uzbekistan Research Online. It has been accepted for inclusion in Scientific Bulletin. Physical and Mathematical Research by an authorized editor of 2030 Uzbekistan Research Online. For more information, please contact sh.erkinov@edu.uz.

УДК 517.98

MATRITSALAR ALGEBRASIDA 2-LOKAL IKKI TOMONLAMA KO'PAYTIRISHLAR TAVSIFI

F.N. Arzikulov, K.A. Solijanova

Maqolada 2-lokal ikki tomonlama ko'paytirish tushunchasi kiritilgan va o'rganilgan. Har qanday barcha komponentalari musbat bo'lgan matritsalar yordamida aniqlangan Δ ikki lokal ikki tomonlama ko'paytirish uchun barcha $X \in M_2(R)$ matritsalarida $\Delta(X) = AXA$ tenglikni qanoatlantiruvchi $A \in M_2(R)$ matritsa mavjudligi isbotlangan. Shuningdek, agar ixtiyoriy barcha komponentalari musbat funksiya bo'lgan matritsalar yordamida aniqlangan $\Delta: M_2(R) \otimes C[a, b] \rightarrow M_2(R) \otimes C[a, b]$ ikki lokal ikki tomonlama ko'paytirish berilgan bo'lsa, u holda barcha $X \in M_2(R) \otimes C[a, b]$ matritsalar uchun $\Delta(X) = AXA$ tenglikni qanoatlantiruvchi $A \in M_2(R) \otimes C[a, b]$ matritsa mavjudligi isbotlangan.

Kalit so'zlar: Ikki o'lchamli matritsalar algebrasi, 2-lokal ikki tomonlama ko'paytirish, uzluksiz funksiyalar algebrasi.

В данной работе введено и исследовано понятие 2-локального двустороннего умножения. Доказано что, для всякого 2-локального двустороннего умножения Δ , определенного матрицами с положительными компонентами существует матрица $A \in M_2(R)$ такая, что $\Delta(X) = AXA$ для всех $X \in M_2(R)$. А также доказано, что для всякого 2-локального двустороннего умножения $\Delta: M_2(R) \otimes C[a, b] \rightarrow M_2(R) \otimes C[a, b]$, определенного матрицами с положительными компонентами существует матрица $A \in M_2(R) \otimes C[a, b]$ такая, что $\Delta(X) = AXA$ для всех $X \in M_2(R) \otimes C[a, b]$.

Ключевые слова: алгебра двумерных матриц, 2-локальное двустороннее умножение, алгебра непрерывных функций.

Kirish

Maqola matritsalar algebrasida ikki lokal ikki tomonlama ko'paytirish mavzusini o'rganishga bag'ishlangan. Bu tushuncha quyidagicha kiritiladi: $\Delta: M_2(R) \rightarrow M_2(R)$ akslantirish uchun ixtiyoriy $X, Y \in M_2(R)$ olganda, shunday $A \in M_2(R)$ mavjud bo'lsaki, bunda $\Delta(x) = AXA$, $\Delta(y) = AYA$ tenglik bajarilsa, Δ ikki lokal ikki tomonlama ko'paytirish deyiladi.

1997-yilda P.Semrl [1] 2-lokal differensiallash tushunchasini kiritgan va cheksiz o'lchovli H separabl Gilbert fazosi ustida aniqlangan barcha chegaralangan chiziqli operatorlarning $B(H)$ algebrasida har qanday 2-lokal differensiallash differensiallash bo'lishini ko'rsatgan. Keyinroq [2] maqolada chekli o'lchovli H Gilbert fazosida aniqlangan $B(H)$ algebra uchun ham shunday natija olingan [3]. Maqolada esa chekli o'lchovli butunlik halqalari ustida aniqlangan matritsalar halqasida har qanday 2-lokal differensiallash differensiallash bo'lishi ko'rsatilgan. [4] Maqolada mualliflar yangi isbotlash usulini ishlab chiqib, Gilbert fazolari uchun yuqorida aytib o'tilgan [1] va [2] maqolalarning natijalarini umumlashtirishgan. Ya'ni, ular ixtiyoriy olingan (separabellik talab etilmaydi) H Gilbert fazosi ustida aniqlangan barcha chiziqli operatorlar $B(H)$ algebrasida 2-lokal differensiallashlarni o'rganishgan va $B(H)$ ustidagi har qanday 2-lokal differensiallash differensiallash bo'lishini isbotlashgan [5-11]. Maqolalarda mualliflar oldingi natijalarni kengaytirishgan va 2-lokal differensiallashlar bo'yicha bir qator natijalar olishgan.

Maqolada 2-lokal ikki tomonlama ko'paytirish tushunchasi kiritilgan va o'rganilgan. Ikki o'lchovli matritsalar algebrasida barcha komponentalari musbat bo'lgan matritsalar yordamida aniqlangan Δ ikki lokal ikki tomonlama ko'paytirish uchun shunday $A \in M_2(R)$ matritsa mavjudligi isbotlanganki, bunda barcha birlik matritsalar $\Delta(e_{ij}) = Ae_{ij}A$ tenglikni qanoatlantiradi. Ushbu tasdiq maqolada lemma ko'rinishida bayon qilingan. Ushbu lemma yordamida, agar ixtiyoriy barcha komponentalari musbat bo'lgan matritsalar yordamida aniqlangan Δ ikki lokal ikki tomonlama ko'paytirish berilgan bo'lsa, u holda barcha $X \in M_2(R)$ matritsalar uchun $\Delta(X) = AXA$ tenglikni qanoatlantiruvchi $A \in M_2(R)$ matritsa mavjudligi haqidagi teorema isbotlangan. Shuningdek, bu teorema $\Delta: M_2(R) \otimes C[a, b] \rightarrow M_2(R) \otimes C[a, b]$ akslantirish uchun ham o'rinli bo'lishi isbotlangan, ya'ni, agar ixtiyoriy barcha komponentalari musbat bo'lgan matritsalar yordamida aniqlangan Δ ikki lokal ikki tomonlama ko'paytirish berilgan bo'lsa, u holda barcha $X \in M_2(R) \otimes C[a, b]$ matritsalar uchun $\Delta(X) = AXA$ tenglikni qanoatlantiruvchi $A \in M_2(R) \otimes C[a, b]$ matritsa mavjudligi isbotlangan.

MATHEMATICS

2-lokal ikki tomonlama ko'paytirish


Ta'rif. Aytaylik, $M_2(R)$ R haqiqiy sonlar maydoni ustida aniqlangan ikki o'lchovli matritsalar algebrasi bo'lsin. Agar $\Delta: M_2(R) \rightarrow M_2(R)$ akslantirish uchun ixtiyoriy $x, y \in M_2(R)$ matritsalarini olganda, shunday $A \in M_2(R)$ mavjud bo'lsaki, bunda $\Delta(x) = AXA$, $\Delta(y) = AY A$ tengliklar bajarilsa, u holda Δ *ikki lokal ikki tomonlama ko'paytirish* deyiladi.

Ushbu kiritilgan tushuncha bo'yicha quyidagi lemma o'rinalidir. Aytaylik, Δ bu $M_2(R)$ ikki o'lchovli matritsalar algebrasida barcha komponentalari musbat bo'lgan matritsalar yordamida aniqlangan ikki lokal ikki tomonlama ko'paytirish bo'lsin.

Lemma 1. Ikki o'lchovli matritsalar algebrasida shunday $A \in M_2(R)$ matritsa mavjudki, barcha birlik $e_{ij} \in M_2(R)$, $i, j = 1, 2$ matritsalar uchun Δ ikki lokal ikki tomonlama ko'paytirishda $\Delta(e_{ij}) = Ae_{ij}A$ tenglik bajariladi, ya'ni:

$$\begin{aligned}\Delta(e_{11}) &= Ae_{11}A, \Delta(e_{12}) = Ae_{12}A \\ \Delta(e_{21}) &= Ae_{21}A, \Delta(e_{22}) = Ae_{22}A.\end{aligned}$$

Isbot. To'rtala $e_{ij} \in M_2(R)$, $i, j = 1, 2$ birlik matritsalarini juft-jufti bilan olsak, oltita juftlik hosil bo'ladi. Ushbu juftliklarning Δ ikki lokal ikki tomonlama ko'paytirishdagi ko'paytma matritsalarini belgilab olaylik.



U holda quyidagi tengliklar o'rinni bo'ladi.

$$\begin{aligned}\Delta(e_{11}) &= Be_{11}B = Ce_{11}C = Ne_{11}N, \Delta(e_{12}) = Be_{12}B = De_{12}D = Me_{12}M, \\ \Delta(e_{21}) &= Ce_{21}C = Ge_{21}G = Me_{21}M, \Delta(e_{22}) = Ge_{22}G = De_{22}D = Ne_{22}N.\end{aligned}$$

Birlik matritsalar uchun ikki lokal ikki tomonlama ko'paytirishni bittadan matritsa uchun hisoblab chiqamiz.

$$\begin{aligned}\Delta(e_{11}) &= Be_{11}B = \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix} = \begin{pmatrix} b_{11}^2 & b_{11}b_{12} \\ b_{21}b_{11} & b_{21}b_{11} \end{pmatrix} \\ \Delta(e_{12}) &= Be_{12}B = \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix} = \begin{pmatrix} b_{11}b_{21} & b_{11}b_{22} \\ b_{21}^2 & b_{21}b_{22} \end{pmatrix} \\ \Delta(e_{21}) &= Ce_{21}C = \begin{pmatrix} c_{11} & c_{12} \\ c_{21} & c_{22} \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} c_{11} & c_{12} \\ c_{21} & c_{22} \end{pmatrix} = \begin{pmatrix} c_{11}c_{12} & c_{12}^2 \\ c_{11}c_{22} & c_{22}c_{12} \end{pmatrix} \\ \Delta(e_{22}) &= Ge_{22}G = \begin{pmatrix} g_{11} & g_{12} \\ g_{21} & g_{22} \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} g_{11} & g_{12} \\ g_{21} & g_{22} \end{pmatrix} = \begin{pmatrix} g_{12}g_{21} & g_{12}g_{22} \\ g_{21}g_{22} & g_{22}^2 \end{pmatrix}\end{aligned}$$

Ikki lokal ikki tomonlama ko'paytirishni bajarayotganda qolgan ikki matritsalariga ko'paytirib hisoblaganimizda ham, ko'paytmadagi matritsa elementlarining indeksleri bir xil bo'ladi. Shu sababli

$$\begin{aligned}Be_{11}B &= Ce_{11}C = Ne_{11}N, Be_{12}B = De_{12}D = Me_{12}M, \\ Ce_{21}C &= Ge_{21}G = Me_{21}M, Ge_{22}G = De_{22}D = Ne_{22}N\end{aligned}$$

tenglikdan ularning barcha elementlarini tenglab chiqamiz:

$$b_{11}^2 = c_{11}^2 = n_{11}^2, b_{11}b_{12} = c_{11}c_{12} = n_{11}n_{12},$$

$$\begin{aligned}
b_{11}b_{21} &= c_{11}c_{21} = n_{11}n_{21}, b_{12}b_{21} = c_{12}c_{21} = n_{12}n_{21}, \\
b_{11}b_{21} &= d_{11}d_{21} = m_{11}m_{21}, b_{21}^2 = d_{21}^2 = m_{21}^2, \\
b_{21}b_{22} &= d_{21}d_{22} = m_{21}m_{22}, c_{11}c_{12} = g_{11}g_{12} = m_{11}m_{12}, \\
c_{12}^2 &= g_{12}^2 = m_{12}^2, c_{11}c_{22} = g_{11}g_{22} = m_{11}m_{22}, \\
c_{12}c_{22} &= g_{12}g_{22} = m_{12}m_{22}, g_{12}g_{21} = d_{12}d_{21} = n_{12}n_{21}, \\
g_{12}g_{22} &= d_{12}d_{22} = n_{12}n_{22}, g_{22}g_{21} = d_{22}d_{21} = n_{22}n_{21}, \\
g_{22}^2 &= d_{22}^2 = n_{22}^2.
\end{aligned}$$

Qaralayotgan matritsalar faqatgina noldan farqli musbat elementlardan tashkil topgan deb qaraymiz. U holda kvadrat tengliklarning elementlari o'zaro teng bo'ladi. Ya'ni:

$$\begin{aligned}
b_{11} &= c_{11} = n_{11}, b_{21} = d_{21} = m_{21} \\
c_{12} &= g_{12} = m_{12}, g_{22} = d_{22} = n_{22}.
\end{aligned}$$

Bu tengliklarni yuqoridagi tengliklarga qo'llaganimizda, quyidagiga ega bo'lamiz:

$$\begin{aligned}
b_{11} &= c_{11} = d_{11} = g_{11} = n_{11} = m_{11}, b_{12} = c_{12} = d_{12} = g_{12} = n_{12} = m_{12}, \\
b_{21} &= c_{21} = d_{21} = g_{21} = n_{21} = m_{21}, b_{22} = c_{22} = d_{22} = g_{22} = n_{22} = m_{22}
\end{aligned}$$

Bundan $B=C=D=G=N=M$. Demak, barcha matritsalar o'zaro tengligiga erishdik. Isbot yakunlandi.

Teorema 1. Aytaylik Δ bu $M_2(R)$ ikki o'lchovli matritsalar algebrasida barcha komponentalari musbat bo'lgan, ya'ni $A = \{a_{ij}, a_{ij} > 0\}$ bo'lgan, matritsalar yordamida aniqlangan ikki lokal ikki tomonlama ko'paytirish bo'lsin. U holda shunday $A \in M_2(R)$ matritsa mavjudki, bunda ixtiyoriy $X \in M_2(R)$ matritsa uchun $\Delta(X) = AXA$ tenglik bajariladi, Δ ikki tomonlama ko'paytirish bo'ladi.

Isbot. Ixtiyoriy $x \in M_2(R)$ uchun

$$\begin{aligned}
\Delta(x) &= BxB, \Delta(e_{11}) = Be_{11}B, \Delta(x) = CxC, \Delta(e_{12}) = Ce_{12}C \\
\Delta(x) &= DxD, \Delta(e_{21}) = De_{21}D, \Delta(x) = FxF, \Delta(e_{22}) = Fe_{22}F.
\end{aligned}$$

1-Lemmaga ko'ra, shunday A mavjudki,

$$Be_{11}B = Ae_{11}A, Ce_{12}C = Ae_{12}A, De_{21}D = Ae_{21}A, Fe_{22}F = Ae_{22}A$$

tengliklar o'rinli bo'ladi. Bundan quyidagilar kelib chiqadi:

$$\begin{aligned}
&\begin{cases} b_{11}^2 = a_{11}^2 \\ b_{11}b_{12} = a_{11}a_{12} \\ b_{11}b_{21} = a_{11}a_{21} \\ b_{21}b_{12} = a_{21}a_{12} \end{cases} \begin{cases} c_{11}c_{21} = a_{11}a_{21} \\ c_{11}c_{22} = a_{11}a_{22} \\ c_{21}^2 = a_{21}^2 \\ c_{21}c_{22} = a_{21}a_{22} \end{cases} \\
&\begin{cases} d_{11}d_{12} = a_{11}a_{12} \\ d_{12}^2 = a_{12}^2 \\ d_{11}d_{22} = a_{11}a_{22} \\ d_{22}d_{12} = a_{22}a_{12} \end{cases} \begin{cases} f_{21}f_{12} = a_{21}a_{12} \\ f_{12}f_{22} = a_{12}a_{22} \\ f_{21}f_{22} = a_{21}a_{22} \\ f_{22}^2 = a_{22}^2 \end{cases}
\end{aligned}$$

ga ega bo'lamiz. Bundan

$$\begin{aligned}
a_{11} &= b_{11} = c_{11} = d_{11}, a_{12} = b_{12} = f_{12} = d_{12}, \\
a_{21} &= b_{21} = c_{21} = f_{21}, a_{22} = c_{22} = d_{22} = f_{22}
\end{aligned}$$

Qulaylik uchun quyidagi belgilashlarni kiritib olamiz:

$$\begin{aligned}
a_{11} &= a_{11}^2x_{11} + a_{11}a_{12}x_{21} + a_{11}a_{21}x_{12} + a_{12}a_{21}x_{22}, \\
a_{12} &= a_{11}a_{12}x_{11} + a_{12}^2x_{21} + a_{11}a_{22}x_{12} + a_{12}a_{22}x_{22}, \\
a_{21} &= a_{21}a_{11}x_{11} + a_{11}a_{22}x_{21} + a_{21}^2x_{12} + a_{21}a_{22}x_{22}, \\
a_{22} &= a_{21}a_{12}x_{11} + a_{12}a_{22}x_{21} + a_{11}a_{22}x_{12} + a_{22}^2x_{22}, \\
\beta_{11} &= b_{11}^2x_{11} + b_{11}b_{12}x_{21} + b_{11}b_{21}x_{12} + b_{12}b_{21}x_{22}, \\
\gamma_{21} &= c_{21}c_{11}x_{11} + c_{11}c_{22}x_{21} + c_{21}^2x_{12} + c_{21}c_{22}x_{22},
\end{aligned}$$

MATHEMATICS

$$\begin{aligned}\delta_{12} &= d_{11}d_{12}x_{11} + d_{12}^2x_{21} + d_{11}d_{22}x_{12} + d_{12}d_{22}x_{22}, \\ \varepsilon_{22} &= f_{21}f_{12}x_{11} + f_{12}f_{22}x_{21} + f_{11}f_{22}x_{12} + f_{22}^2x_{22}.\end{aligned}$$

U holda

$$\begin{aligned}\Delta(x) &= BXB = \begin{pmatrix} \beta_{11} & \alpha_{12} \\ \alpha_{21} & \alpha_{22} \end{pmatrix}, \Delta(x) = CXC = \begin{pmatrix} \alpha_{11} & \alpha_{12} \\ \gamma_{21} & \alpha_{22} \end{pmatrix}, \\ \Delta(x) &= DXD = \begin{pmatrix} \alpha_{11} & \delta_{12} \\ \alpha_{21} & \alpha_{22} \end{pmatrix}, \Delta(x) = FXF = \begin{pmatrix} \alpha_{11} & \alpha_{12} \\ \alpha_{21} & \varepsilon_{22} \end{pmatrix}.\end{aligned}$$

Demak,

$$\Delta(x) = BXB = CXC = DXD = FXF = \begin{pmatrix} \alpha_{11} & \alpha_{12} \\ \alpha_{21} & \alpha_{22} \end{pmatrix}.$$

Shunday $A \in M_2(R)$ borki, ixtiyoriy $x \in M_2(R)$ olinganda, $\Delta(X) = AXA$ tenglik o`rinli bo`ladi. Teorema isbotlandi.

Aytaylik, $M_2(R) \otimes C[a, b]$ bu komponentalari $C[a, b]$ uzluksiz funksiyalar algebrasidan olingan barcha 2×2 o`lchovli matritsalar algebrasi bo`lsin.

Lemma 3. Aytaylik, Δ bu $M_2(R) \otimes C[a, b]$ ikki o`lchovli matritsalar algebrasida barcha komponentalari musbat funksiya bo`lgan matritsalar yordamida aniqlangan ikki lokal ikki tomonlama ko`paytirish bo`lsin. U holda ikki o`lchovli matritsalar algebrasida shunday $A \in M_2(R) \otimes C[a, b]$ matritsa mavjudki, barcha birlik $e_{ij}(t) \in M_2(R) \otimes C[a, b]$, $i, j = 1, 2$ matritsalar uchun Δ ikki lokal ikki tomonlama ko`paytirishda $\Delta(e_{ij}) = Ae_{ij}A$ tenglik bajariladi, ya`ni:

$$\begin{aligned}\Delta(e_{11}(t)) &= Ae_{11}(t)A, \Delta(e_{12}(t)) = Ae_{12}(t)A \\ \Delta(e_{21}(t)) &= Ae_{21}(t)A, \Delta(e_{22}(t)) = Ae_{22}(t)A.\end{aligned}$$

Teorema 4. Aytaylik, Δ bu $M_2(R) \otimes C[a, b]$ ikki o`lchovli matritsalar algebrasida barcha komponentalari musbat funksiya bo`lgan matritsalar yordamida aniqlangan ikki lokal ikki tomonlama ko`paytirish bo`lsin, ya`ni $\Delta: M_2(R) \otimes C[a, b] \rightarrow M_2(R) \otimes C[a, b]$ shunday ikki lokal ikki tomonlama ko`paytirish bo`lsinki ixtiyoriy $x, y \in M_2(R) \otimes C[a, b]$ elementlar uchun, $\Delta(x) = axa$ $\Delta(y) = aya$ shartni qanoatlantiruvchi $a = \begin{pmatrix} a_{11}(t) & a_{12}(t) \\ a_{21}(t) & a_{22}(t) \end{pmatrix}$ matritsa uchun, ixtiyoriy $x \in [a, b]$ uchun $a_{11}(t) > 0, a_{12}(t) > 0, a_{21}(t) > 0, a_{22}(t) > 0$ bo`lsin. U holda Δ ikki tomonlama ko`paytirish bo`ladi, ya`ni shunday $a \in M_2(R) \otimes C[a, b]$ topiladiki, ixtiyoriy $x \in M_2(R) \otimes C[a, b]$ uchun $\Delta(x) = axa$ o`rinli bo`ladi.

Isbot. Ixtiyoriy $x \in M_2(R) \otimes C[a, b]$ uchun

$$\begin{aligned}\Delta(x) &= BxB, \Delta(e_{11}(t)) = Be_{11}(t)B \\ \Delta(x) &= CxC, \Delta(e_{12}(t)) = Ce_{12}(t)C \\ \Delta(x) &= DxD, \Delta(e_{21}(t)) = De_{21}(t)D \\ \Delta(x) &= FxF, \Delta(e_{22}(t)) = Fe_{22}(t)F.\end{aligned}$$

3-Lemmaga ko`ra, shunday A mavjudki,

$$\begin{aligned}Be_{11}(t)B &= Ae_{11}(t)A, Ce_{12}(t)C = Ae_{12}(t)A, \\ De_{21}(t)D &= Ae_{21}(t)A, Fe_{22}(t)F = Ae_{22}(t)A\end{aligned}$$

tenglik o`rinli, ya`ni:

$$\begin{cases} b_{11}^2(t) = a_{11}^2(t) \\ b_{11}(t)b_{12}(t) = a_{11}(t)a_{12}(t) \\ b_{11}(t)b_{21}(t) = a_{11}(t)a_{21}(t) \\ b_{21}(t)b_{12}(t) = a_{21}(t)a_{12}(t) \end{cases}$$

$$\begin{cases} c_{11}(t)c_{21}(t) = a_{11}(t)a_{21}(t) \\ c_{11}(t)c_{22}(t) = a_{11}(t)a_{22}(t) \\ c_{21}^2(t) = a_{21}^2(t) \\ c_{21}(t)c_{22}(t) = a_{21}(t)a_{22}(t) \end{cases}$$

$$\begin{cases} d_{11}(t)d_{12}(t) = a_{11}(t)a_{12}(t) \\ d_{12}^2(t) = a_{12}^2(t) \\ d_{11}(t)d_{22}(t) = a_{11}(t)a_{22}(t) \\ d_{22}(t)d_{12}(t) = a_{22}(t)a_{12}(t) \end{cases}$$

$$\begin{cases} f_{21}(t)f_{12}(t) = a_{21}(t)a_{12}(t) \\ f_{12}(t)f_{22}(t) = a_{12}(t)a_{22}(t) \\ f_{21}(t)f_{22}(t) = a_{21}(t)a_{22}(t) \\ f_{22}^2(t) = a_{22}^2(t) \end{cases}$$

ga ega bo'lamiz. Bundan

$$\begin{aligned} a_{11}(t) &= b_{11}(t) = c_{11}(t) = d_{11}(t), a_{12}(t) = b_{12}(t) = f_{12}(t) = d_{12}(t), \\ a_{21}(t) &= b_{21}(t) = c_{21}(t) = f_{21}(t), a_{22}(t) = c_{22}(t) = d_{22}(t) = f_{22}(t) \end{aligned}$$

Qulaylik uchun quyidagi belgilashlarni kiritib olamiz:

$$\begin{aligned} \alpha_{11}(t) &= a_{11}^2(t)x_{11} + a_{11}(t)a_{12}(t)x_{21} + a_{11}(t)a_{21}(t)x_{12} + a_{12}(t)a_{21}(t)x_{22}, \\ \alpha_{12} &= a_{11}(t)a_{12}(t)x_{11} + a_{12}^2(t)x_{21} + a_{11}(t)a_{22}(t)x_{12} + a_{12}(t)a_{22}(t)x_{22}, \\ \alpha_{21}(t) &= a_{21}(t)a_{11}(t)x_{11} + a_{11}(t)a_{22}(t)x_{21} + a_{21}^2(t)x_{12} + a_{21}(t)a_{22}(t)x_{22}, \\ \alpha_{22}(t) &= a_{21}(t)a_{12}(t)x_{11} + a_{12}(t)a_{22}(t)x_{21} + a_{11}(t)a_{22}(t)x_{12} + a_{22}^2(t)x_{22}, \\ \beta_{11}(t) &= b_{11}^2(t)x_{11} + b_{11}(t)b_{12}(t)x_{21} + b_{11}(t)b_{21}(t)x_{12} + b_{12}(t)b_{21}(t)x_{22}, \\ \gamma_{21}(t) &= c_{21}(t)c_{11}(t)x_{11} + c_{11}(t)c_{22}(t)x_{21} + c_{21}^2(t)x_{12} + c_{21}(t)c_{22}(t)x_{22}, \\ \delta_{12}(t) &= d_{11}(t)d_{12}(t)x_{11} + d_{12}^2(t)x_{21} + d_{11}(t)d_{22}(t)x_{12} + d_{12}(t)d_{22}(t)x_{22}, \\ \varepsilon_{22}(t) &= f_{21}(t)f_{12}(t)x_{11} + f_{12}(t)f_{22}(t)x_{21} + f_{11}(t)f_{22}(t)x_{12} + f_{22}^2(t)x_{22}. \end{aligned}$$

U holda

$$\begin{aligned} \Delta(x) &= BXB = \begin{pmatrix} \beta_{11}(t) & \alpha_{12}(t) \\ \alpha_{21}(t) & \alpha_{22}(t) \end{pmatrix}, \Delta(x) = CXC = \begin{pmatrix} \alpha_{11}(t) & \alpha_{12}(t) \\ \gamma_{21}(t) & \alpha_{22}(t) \end{pmatrix}, \\ \Delta(x) &= DXD = \begin{pmatrix} \alpha_{11}(t) & \delta_{12}(t) \\ \alpha_{21}(t) & \alpha_{22}(t) \end{pmatrix}, \Delta(x) = FXF = \begin{pmatrix} \alpha_{11}(t) & \alpha_{12}(t) \\ \alpha_{21}(t) & \varepsilon_{22}(t) \end{pmatrix}. \end{aligned}$$

Demak,

$$\Delta(x) = BXB = CXC = DXD = FXF = \begin{pmatrix} \alpha_{11}(t) & \alpha_{12}(t) \\ \alpha_{21}(t) & \alpha_{22}(t) \end{pmatrix}.$$

Teorema isbotlandi.

References

1. Šemrl, P. (1997) Lokal automorphisms and derivations on $B(H)$. *Proceedings of the American Mathematical Society*. 125(9). Pp. 2677-2680.
2. Kim S., Kim J. (2004) Lokal automorphisms and derivations on M_n . *Proceedings of the American Mathematical Society*. 132 1389-1392.
3. Lin Y., Wong T. (2006) A note on 2-lokal maps. *Proceedings of the Edinburgh Mathematical Society*. 49. Pp. 701-708.
4. Ayupov Sh., Kudaybergenov K. (2012) 2-lokal derivations and automorphisms on $B(H)$. *Journal of Mathematical Analysis and Applications*. 395. Pp. 15-18.
5. Ayupov Sh., Arzikulov F. (2014) 2-Lokal derivations on semi-finite von Neumann algebras, *Glasgow Mathematical Journal*. 56. Pp. 9-12.
6. Ayupov Sh., Kudaybergenov K. (2015) 2-Lokal derivations on von Neumann algebras. *Positivity*. 19. Pp. 445-455.
7. Ayupov S., Arzikulov F. (2017) 2-Lokal derivations on associative and Jordan matrix rings over commutative rings. *Linear Algebra and its Applications*. 522. Pp. 28-50.
8. Ayupov Sh.A., Arzikulov F.N. (2018) Description of 2-lokal and lokal derivations on some Lie rings of skew-adjoint matrices. *Linear and Multilinear Algebra*. 9(26). Pp. 31-17
9. Ayupov Sh.A., Arzikulov F.N. (2017) 2-Lokal Derivations on AW^* -Algebras of Type I. *Lobachevskii Journal of Mathematics*. 1(38). Pp. 148-161.

MATHEMATICS

10. Ayupov Sh.A., Arzikulov F.N. (2018) 2-Lokal derivations on algebras of matrix-valued functions on a compact. *Vladikavkazskiy matematicheskiy jurnal*. Vol. 20. Issue 1. Pp. 38-49.

11. Ayupov Sh.A., Arzikulov F.N., Umrzaqov N.M. (2019) Lokal derivations on associative and Jordan matrix algebras. *O'zbek matematika jurnali*. 4. Pp. 13-33.

DESCRIPTION OF 2-LOCAL TWO SIDED MULTIPLICATION ON AN ALGEBRA OF MATRICES

F.N. Arzikulov¹, K.A. Solijanov¹

Ilmiy xabarnoma. Fizika-matematika tadqiqotlari – Scientific Bulletin. Physical and Mathematical Research. 2019. 2. 73–78.

¹Andijan State University, Andijan, 170100, str. University, 129 (Uzbekistan). E-mail: agsu_info@edu.uz

Key words: algebra of matrices on R , two sided multiplication, 2-local two sided multiplication

The present paper is devoted to 2-local derivation on associative and Jordan matrix rings. In 1997, P.Semrl introduced the notion of 2-local derivations and described 2-local derivations on the algebra $B(H)$ of all bounded linear operators on the infinite-dimensional separable Hilbert space H . A similar description for the finite-dimensional case appeared later in 2004. In the paper Y. Lin and T. Wong 2-local derivations have been described on matrix algebras over finite dimensional division rings. In 2012 Sh. Ayupov, K. Kudaybergenov suggested a new technique and generalized the above mentioned results for arbitrary Hilbert spaces. Namely they considered 2-local derivations on the algebra $B(H)$ of all linear bounded operators on an arbitrary (no separability is assumed) Hilbert space H and proved that every 2-local derivation on $B(H)$ is a derivation. Later Sh.Ayupov, K.Kudaybergenov and F.Arizkulov extended the above results and gave a proof of the theorem for an arbitrary von Neumann algebra.

In the given paper it is introduced and studied a concept of 2-local two sided multiplication on $M_2(R)$. A 2-local two sided multiplication is defined as follows: let $\Delta: M_2R \rightarrow M_2R$ be a mapping. If for each pairs $X, Y \in M_2(R)$ of elements there exists a matrix $A \in M_2(R)$ such that $\Delta(x) = AXA$, $\Delta(y) = AYA$, then the mapping Δ is called 2-local two-sided multiplication. It is proved that every 2-local two sided multiplication of $M_2(R)$ is a two sided multiplication, if all components of all matrices defining this 2-local two sided multiplication are positive. Namely, we prove that for any matrix $x \in M_2(R)$ there exists a matrix $a \in M_2(R)$ with positive entries such that $\varphi(x) = axa$. We prove the following lemma: let Δ be a 2-local two sided multiplication on $M_2(R)$ defined by 2×2 matrices with positive entries of the field of real numbers. Then there exists a matrix $A \in M_2(R)$ with positive entries such that, for all matrix units $e_{ij} \in M_2(R)$, $i, j = 1, 2$ it is valid $\Delta(e_{ij}) = Ae_{ij}A$ for the 2-local two sided multiplication Δ , that is $\Delta(e_{11}) = Ae_{11}A$, $\Delta(e_{12}) = Ae_{12}A$, $\Delta(e_{21}) = Ae_{21}A$, $\Delta(e_{22}) = Ae_{22}A$.

Similarly we define a concept of 2-local two sided multiplication on $M_2(R) \otimes C[a, b]$ as follows: let $\Delta: M_2R \otimes C[a, b] \rightarrow M_2R \otimes C[a, b]$ be a mapping. If for each pairs $X, Y \in M_2(R) \otimes C[a, b]$ of elements there exists a matrix $A \in M_2(R) \otimes C[a, b]$ such that $\Delta(x) = AXA$, $\Delta(y) = AYA$, then the mapping Δ is called 2-local two-sided multiplication. In the present paper, it is proved that every 2-local two sided multiplication of $M_2(R) \otimes C[a, b]$ is a two sided multiplication, if all components of all matrices defining this 2-local two sided multiplication are positive functions. Namely, we prove that for any matrix $x \in M_2(R) \otimes C[a, b]$ there exists a matrix $a \in M_2(R)$ with positive function entries such that $\varphi(x) = axa$.

Муаллифлар ҳақида маълумот:

Арзикулов Фарходжон Нематжонович – физика-математика фанлари доктори, Андижон давлат университети математика кафедраси доценти. E-mail: arzikulovfn@rambler.ru

Солижанова Камола Абдушукур қизи – Андижон давлат университети математика кафедраси магистранти.